

Expression & purification of 6his-SmPRMT3(1-553)

Construct

MGSSHHHHHHSSGTGSGENLYFQGHMSPCDQSSNYDLDMEPNDSSYFGSYGHFEIHGEMINDRVRTESYVNFILSN
AEKYFKHKIILDVGS GSGILSIIAAQAGASHVYGVEAADEIYAASHETLRVNNLLERVTFIHGQAESVELPVKKVDVIISE
WMGYFLFFESMLDSVLKMASKYLSRDGHIFPRHYTLNLLGVQCSEQLRKRRLEHWNNVYGYNMPALRRAALSEAHV
LNLTNEHVTPPISPITILTQSFELVALDLDDMHRNRINYLSNHCSLLCEQKFHLTIQPTTDINNNSSSSSSYELDAIVGYFD
VRFDDADCKVEFSTSPPTPLTHWKQTLFLDKPIRVKPGDKISGIITIRRATTDNRGLEINLLIGETENSLEIKQTFDLIG

Length = 396 AA

Molecular weight = 44782.3 Da

pI 5.66

$\xi(\text{red})$ 40340 L.M⁻¹

Expresion

Grow pnEAvH_SmPRMT3-184 BL21(DE3) transformants on an LB-Amp plate (1 plate/L culture), overnight 37°C. Resuspend cells in LB medium (5 mL/plate) and inoculate liquide **2X** LB medium + 100 µg/mL⁻¹ Ampicilline. Grow culture(s) at 37°C, 200 rpm and measure OD600 every 20 minutes. When OD600 reaches 0.4, set temperature to 20°C. When OD600 reaches 0.6 to 0.8, induce 6his-SmPRMT3(184-553) with 1 mM IPTG.

From:

<https://bsi.inscog.eu/> - BSI wiki

Permanent link:

https://bsi.inscog.eu/doku.php?id=purification:expression_and_purification_of_smprmt3_184-553&rev=1477661600

Last update: 2023/11/01 20:17

